

# Recent CLEO Results

Karl M. Ecklund

Cornell University

Beauty 2003


October 14, 2003

Carnegie Mellon University

Web site: [www-hep.phys.cmu.edu/beauty2003/](http://www-hep.phys.cmu.edu/beauty2003/)  
Email: [beauty2003@phys.cmu.edu](mailto:beauty2003@phys.cmu.edu)

Carnegie Mellon


# CLEO CKM Results



- Motivation
  - Physics of flavor
  - CPV from CKM?
  - Look for new physics
- CLEO's contribution:  $|V_{ub}|, |V_{cb}|$  from semileptonic B decays
  - Pioneering measurements
  - Still among the best!

Unitarity Triangle

# UT Constraint from $|V_{ub}|$



$|V_{ub}|$  from  $B \rightarrow \pi \ell \nu$ :


$$\frac{d\Gamma}{dq^2} = \frac{G_F^2}{24\pi^3} |V_{ub}|^2 p_\pi^3 |f_+(q^2)|^2$$

Form factor  $f(q^2)$ :

- Encodes hadronic physics
- Not well known
- Limits  $|V_{ub}|$  precision

- CLEO has measured  $B \rightarrow \pi \ell \nu$ ,  $\rho \ell \nu$  before
- New measurement that is binned in  $q^2$ , therefore sensitive to shape of  $f(q^2)$
- FF computed in quark models, LQCD and LCSR

# Exclusive $B \rightarrow \pi \ell \nu$




- Suppress  $b \rightarrow c$  bkgd by reconstructing  $\nu$
- Use hermeticity of detector to infer  $p_\nu$
- Clean events required
  - Remove spurious tracks
  - and hadronic showers
- $(E, p)$  conservation  $\rightarrow$  peaks in  $M_B$  &  $\Delta E$
- Rate and form factor give  $|V_{ub}|$

# Exclusive $|V_{ub}|$

- 7  $B \rightarrow X_{\ell} \ell \nu$  submodes considered ( $\pi, \rho, \omega, \eta$ )
- 3  $q^2$  bins for  $\pi, \rho$
- Simultaneous ML Fit
  - Accounts for cross feed
  - Fit projections shown on right
- Isospin constraints
  - $\frac{1}{2}\Gamma(\pi^- \ell \nu) = \Gamma(\pi^0 \ell \nu)$
  - $\frac{1}{2}\Gamma(\rho^- \ell \nu) = \Gamma(\rho^0 \ell \nu) \approx \Gamma(\omega \ell \nu)$

$$BF(B^0 \rightarrow \pi^- \ell \nu) = (1.33 \pm 0.18 \pm 0.11 \pm 0.01 \pm 0.07) \times 10^{-4}$$



$$BF(B^0 \rightarrow \rho^- \ell \nu) = (2.17 \pm 0.34^{+0.47}_{-0.54} \pm 0.01 \pm 0.41) \times 10^{-4}$$



# Extracting $|V_{ub}|$

$$\frac{d\Gamma(B^0 \rightarrow P^- \ell^+ \nu)}{dy d\cos\theta_{W\ell}} = |V_{ub}|^2 \frac{G_F^2 k_P^3 M_B^2}{32\pi^3} \sin^2\theta_{W\ell} |f_1(q^2)|^2$$



- Fit  $d\Gamma/dq^2$ 
  - Discriminates among FFs
- $B \rightarrow \pi \ell \nu$ 
  - FF dependence is small
  - Disfavors ISGW2
- $B \rightarrow \rho \ell \nu$ 
  - Larger FF dependence  $\Rightarrow$  greater model uncertainty in  $|V_{ub}|$



# Results

$B \rightarrow \pi l \nu$


$B \rightarrow \rho l \nu$



PRD 68, 072003 (2003)

$$|V_{ub}| = (3.17 \pm 0.17 \pm 0.17 \begin{smallmatrix} +0.53 \\ -0.39 \end{smallmatrix} \pm 0.03) \times 10^{-3}$$

# World $|V_{ub}|$ Results




- All measurements are systematics limited
- CLEO pioneering new techniques and
- Using a very well-understood detector
- CLEO results still very competitive in B factory era

Ed Thorndike's Compilation @ FPCP `03



# B Semileptonic Branching Fraction


- CLEO II:  $10 \text{ fb}^{-1}$  at  $\Upsilon(4S)$ 
  - Mature, well-understood detector, data, Monte Carlo, generators, etc.
- Lepton-Tagged Analysis
  - $p_{\text{tag}} > 1.4 \text{ GeV}/c$  plus accompanying electron with  $p_e > 0.6 \text{ GeV}/c$ .
  - Charge, angular correlations to separate primary ( $B \rightarrow X e \nu$ ) from secondary ( $B \rightarrow D \rightarrow Y e \nu$ ).
- Refined electron ID, background and efficiency determinations.
  - Maximize understanding and minimize momentum dependence.



$$\mathcal{B}(B \rightarrow X e \nu) = (10.88 \pm 0.08 \pm 0.33)\% \quad \text{stat} \pm \text{syst}$$

# $B_{SL}$ - Status

- Good agreement among different techniques, experiments.
- Measurements at  $\Upsilon(4S)$  have come up and LEP  $Z^0$  average has come down.
  - Most recent LEP fit result is  $(10.59 \pm 0.22)\%$



# Inclusive $|V_{cb}|$

Heavy Quark Expansion: double series in  $1/M, \alpha_s$

$$\Gamma_{sl} = \frac{G_F^2 |V_{cb}|^2 M_B^5}{192\pi^3} 0.3689 \left[ 1 - 1.54 \frac{\alpha_s}{\pi} - 1.43 \beta_0 \frac{\alpha_s^2}{\pi^2} - 1.648 \frac{\bar{\Lambda}}{M_B} (1 - 0.87 \frac{\alpha_s}{\pi}) - 0.946 \frac{\bar{\Lambda}^2}{M_B^2} - 3.185 \frac{\lambda_1}{M_B^2} \right. \\ \left. + 0.02 \frac{\lambda_2}{M_B^2} - 0.298 \frac{\bar{\Lambda}^3}{M_B^3} - 3.28 \frac{\bar{\Lambda} \lambda_1}{M_B^3} + 10.47 \frac{\bar{\Lambda} \lambda_2}{M_B^3} - 6.153 \frac{\rho_1}{M_B^3} + 7.482 \frac{\rho_2}{M_B^3} \right. \\ \left. - 7.4 \frac{T_1}{M_B^3} + 1.491 \frac{T_2}{M_B^3} - 10.41 \frac{T_3}{M_B^3} - 7.482 \frac{T_4}{M_B^3} + \mathcal{O}(1/M_B^4) \right].$$

- **Ingredients:**

- $B(B \rightarrow X_{\perp} \nu) = (10.8 \pm 0.3)\%$  (CLEO)
- $\tau_{B^0}$  and  $\tau_{B^\pm}$  (PDG),  $f_{+/-} / f_{00}$  (CLEO)
- HQE parameters  $\Lambda, \lambda_1$ , from moments  $\langle E_\gamma \rangle: B \rightarrow X_s \gamma, \langle M_X^2 \rangle: B \rightarrow X_{\perp} \nu$  (CLEO)
- HQE parameter  $\lambda_2 = 0.128 \pm 0.010$  from  $B^* - B$  mass difference  
 $\Rightarrow \Gamma_{sl} = (0.44 \pm 0.02) \times 10^{-10} \text{ MeV}$

- **Result:**

$$|V_{cb}| = 0.0411 \pm 0.0005_{\text{exp } \Lambda, \lambda_1} \pm 0.0007_{\text{exp } \Gamma} \pm 0.0009_{\text{theory}}$$


Overall precision:  $\sim 3\%$  + quark-hadron duality.

# New and Improved Measurement of the Hadronic Mass Moments in $B \rightarrow X_c \ell \nu$

hep-ex/0307081 contributed to Lepton-Photon 2003

- Compute recoiling hadronic mass from charged lepton and neutrino kinematics - neutrino "reconstruction"
- Near hermeticity of CLEO II  $\Rightarrow$  Neutrino="What's missing"

Preliminary




Fit 3-dimensional differential decay rate, extract hadronic mass squared as a function of lepton-energy cut ( $p_\ell > 1 \text{ GeV}/c$ ).

$$M_X^2 = M_B^2 + q^2 - 2E_{beam}(E_\ell + E_\nu) + \underbrace{2|\vec{p}_B||\vec{q}|\cos\theta_{B-q}}_{\text{neglected}}$$

# Fit Projections

- Selection criteria:
  - Cuts to enhance  $\nu$  reconstruction
  - Continuum suppression
  - Efficiency  $\sim 2\%$  for  $B \rightarrow X_c l \nu$
- Sample to fit: 122K events
- Components of fit:
  - $B \rightarrow D l \nu$  } HQET+measured FFs
  - $B \rightarrow D^* l \nu$  }
  - $B \rightarrow D^{**} l \nu$  ISGW2
  - $B \rightarrow (X_c)_{NR} l \nu$  Goity/Roberts
  - $B \rightarrow X_u l \nu$  ISGW2+NR
  - Secondaries CLEO MC
  - Fake Leptons, Continuum fixed with data



# Results

Fits → Mode-by-Mode BFs


+

Generator-Level Info  
(Fraction above  $p_1$  cut, moment  
value for that cut.)



Moment Value for  $B \rightarrow X_c \ell \nu$

hep-ex/0307081



Cut (GeV)	$\langle M_X^2 - \bar{M}_D^2 \rangle$ ( $\text{GeV}^2/c^4$ )
$E_\ell > 1.0$	$0.456 \pm 0.014 \pm 0.045 \pm 0.109$
$E_\ell > 1.1$	$0.422 \pm 0.014 \pm 0.031 \pm 0.084$
$E_\ell > 1.2$	$0.393 \pm 0.013 \pm 0.027 \pm 0.069$
$E_\ell > 1.3$	$0.364 \pm 0.013 \pm 0.030 \pm 0.054$
$E_\ell > 1.4$	$0.332 \pm 0.012 \pm 0.027 \pm 0.055$
$E_\ell > 1.5$	$0.293 \pm 0.012 \pm 0.033 \pm 0.048$

stat  $\pm$  syst  $\pm$  model

- Consistent with previous CLEO measurements, BaBar summer '03, DELPHI
- Interpretation is ongoing

# More CLEO Physics Results

- Rare B decays
  - $A_{CP}$  in  $B \rightarrow K^{*+}\pi^-$
  - $B \rightarrow \eta' X_s$  BF
  - Upper Limit on Baryons in  $B \rightarrow X_s \gamma$
- Hadronic B Decays
  - $B \rightarrow D^{(*)} \rho$  helicity analysis (Final State Interactions)
- Upsilon Decays
  - $Y(3S) \rightarrow \omega Y(1S)$
  - Two-body  $Y(nS)$  decays
  - Searches for  $cc$  states
- Charmed Baryons
  - CPV in  $\Lambda_c \rightarrow \Lambda e \nu$
- Charm Decays
  - Branching fractions
  - Mixing and DCSD
  - Dalitz plot analyses
    - Hadronic structure
    - CPV via interference in Dalitz Plot  $D^0 \rightarrow \pi^+ \pi^- \pi^0$
- $D_s$  spectroscopy
  - See talk by JC Wang

# Summary

- CLEO is still contributing to  $|V_{ub}|$  and  $|V_{cb}|$  measurements
  - $|V_{cb}|$
  - $|V_{ub}|$
- See talks by
  - David Asner on CLEO-c Prospects
  - J.C. Wang on the new  $D_{sJ}$  states